

Insights into Structural Analysis via Crystallographic and X-ray Diffraction Techniques

M Kishore Babu¹, P.Swathi², Md.Yunoos³, R.V.S.Mounica⁴

1. Professor, Department of Pharmaceutics, QIS College of pharmacy, Ongole, A.P
2. Associate Professor, Department of pharmaceutical chemistry, QIS College of pharmacy, Ongole, A.P
3. Professor, Department of pharmaceutical Analysis , QIS College of pharmacy, Ongole, A.P
4. Assistant Professor, Department of pharmaceutical Analysis , QIS College of pharmacy, Ongole, A.P

To Cite this Article

M Kishore Babu, P.Swathi, Md.Yunoos, R.V.S.Mounica, "Insights into Structural Analysis via Crystallographic and X-ray Diffraction Techniques" *Journal of Science and Technology*, Vol. 10, Issue 01- Jan 2025, pp38-47

Article Info

Received: 29-12-2024 Revised: 06-01-2025 Accepted: 17-01-2025 Published: 27-01-2025

Abstract

The arrangement of molecules inside a solid directly affects its properties. Therefore, understanding the three-dimensional structure of matter is necessary for establishing relationships between arrangement and qualities as well as for developing useful materials. X-ray crystallography has been the preferred method for precisely characterizing molecule structure at the atomic level for the last century. Crystallography is a fundamental analytical method that is often helpful in the development of novel medications. Atom size, length, electron density, atom mean location, and other attributes are among its attributes. The method is accurate for analyzing and determining the molecule and molecular configuration of crystals. X-ray diffraction (XRD) is the process by which an x-ray beam disperses or scatters into many directions and takes on unique patterns or shapes due to the presence of crystalline particles. Numerous industries, including the pharmaceutical sector, nanotechnology, geology, electronics, mineralogy, geosciences, materials science, and medicine, benefit from the use of X-ray diffraction and the crystallography method. Because of the structural knowledge gained from crystallographic research, these and other areas have developed quickly.

Keywords: Bragg's law, Crystallography, Diffraction, Diffraction methods, X-ray

1. The Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which permits others to remix, modify, and expand upon the work noncommercially as long as proper credit is provided and the new creations are licensed under the same terms, governs the distribution of articles in this Open Access (OA) journal.

Corresponding Author: M. Kishore Babu

Mail: kishorebabu.m@gmail.com

2. Introduction

2.1. X ray

3. X-rays are electromagnetic radiations (EMR) that have wavelengths between 10-3 and 10 nm. They were given that moniker because, when W. Rontgen discovered X-rays in 1895, their qualities were unknown. One X-rays are far more energetic and have shorter wavelengths than UV light. These are helpful for taking pictures of the human body since they can see through a person's skin to the bones underneath it. One Soft X-rays have longer wavelengths than hard X-rays, which have wavelengths between 0.1 and 0.2 nm. The kind of X-rays utilized in material analysis are called hard X-rays. 2.

3.1. Diffraction phenomenon

The diffraction of light had already been known before X- rays were discovered. The term "diffraction" describes a number of phenomena that happen when a wave runs into a slit or an obstruction. The diffraction phenomena are

defined in classical physics as the spreading out of waves flowing through small holes and the bending of waves around small obstructions.²

4. X-ray Crystallography

The major technique for accessing the arrangement of the molecule or atom is X-ray crystallography. The three-dimensional structure and x-ray crystallography procedures are accurate, providing a three-dimensional structural parameter that is important for drug design and enabling functional research that are structurally based.³

A concentrated amount of a refined substance crystallizes, and the crystals are subsequently exposed to an x-ray beam. Subsequently, the diffraction patterns that have been processed can be utilized to obtain more information regarding the symmetry of the crystal packing and the dimensions of its repeating unit.⁴

4.1. X-ray diffraction

One technique that is non-invasive is X-ray diffraction (XRD) also aids in understanding how atoms are arranged, how these configurations can affect the behavior of materials, or how they can be utilized to ascertain the structure of complicated compounds.⁵ Additionally, XRD is helpful in evaluating the qualitative and quantitative identification of different chemical compounds, assessing the intensity or degree of crystallinity of compounds, and identifying stacking faults (it is a type of defect). The examination of different compound features, such as particle size, polymorphism, isomorphism, phase transition, etc., is frequently done using XRD.³ XRD has various types and one of commonly used is PXRD / XRPD. X-Ray Powder Diffraction (XRPD) provides firsthand knowledge into the elements and their physical and chemical characteristics.⁶

5. Background of X-ray diffraction and crystallography

Numerous groundbreaking works led to the development of X-ray crystallography as a technique for precise solid-state substance analysis. W.C. Roentgen made the discovery of X- rays in 1895. M. von Laue proved in 1912 that single crystals may diffract X-rays, and for this work the Nobel Prize in Physics was given to him in 1914 (Friedrich W, Knipping P, Laue M, 1912). The diffraction law known as Bragg's law was developed by W.L. Bragg, who also demonstrated how to use a diffraction pattern to determine the crystal structure of NaCl.⁷ Instruments for observing diffraction patterns were first designed by his father, W. H. Bragg, who was a pioneer in the field. The diffraction pattern and their combined and individual diffraction patterns were analyzed using Bragg's law. The father and son Braggs received the 1915 Physics Nobel Prize as a result of this.⁸

5.1. Principle of X-ray diffraction crystallography

A unidirectional X-ray beam focuses on the object being examined in order to resolve details of structure within the lattice of the crystal. The materials typically consist of crystallized repeating homogeneous atomic planes. Typically, a specific tube called a cathode ray tube is used to produce polychromatic X-rays. When polychromatic X-rays are passed via a monochromator, monochromatic radiation is created, which distinguishes the scattered, transferred, & absorbing rays upon reaching their atomic structures.⁹ In a sealed tube with a vacuum atmosphere, X-rays are created. When 15–60 kilovolts of current are applied inside the tube, electrons are produced that hit an anode made of Cr, Fe, Co, Cu, Mo, or Ag and produce X-ray beams.

Bragg's law states that incoming X-rays react with the specimen's atomic layers to produce diffracted, transmitted, refracted, scattered, and absorbed beam.

$n\lambda = 2d \sin\theta$ Where,

λ - Wavelength of the ray

θ – Angle between incident rays and surface of crystal. d – Spacing between layers of atoms

Constructive interference occurs when n is integer / whole no.

This equation dealt with the relationship between electromagnetic radiation, the crystal sample's lattice spacing (the physical size of a unit cell in a crystallattice), and the diffraction angle. We have identified, analyzed, and counted these diffracted x-rays. Using a range 2θ angle to scan the sample. Lattice diffraction should occur in every direction that is practical. Since every mineral has a unique set of d spacing, changing the diffraction peak to d-space enables mineral identification. Typically, a difference of d spacing with a standard reference sample is used to achieve it.³

5.2. Bragg's law of diffraction

W.L. Bragg thought of X-ray diffraction through the surface of a crystal as an obstacle of reflecting X-rays across the atomic structures of the minerals in line with the principles of reflection with the objective to comprehend the diffraction of X-rays.

When X-rays that have been scattered, react constructively with one another at certain wavelengths with particular incidence angles, they form powerful reflecting X-rays. In order for constructive interference to occur, the differences in the electromagnetic paths must be an integer number of the wavelength. When this constructive interference occurs, a scattered X-ray beam escapes the crystal at an angle corresponding with the angle of the incident beam.¹¹

Think of a collection of perpendicular planes of atoms in the crystal that have Miller indices and a spacing of d among each succeeding surface. Assume the incident rays of a straight beam of X-rays that are monochromatic of wavelength λ lay in the horizontal direction of the substrate at a striking point θ .




Figure 1: Explanation of bragg's law

Allow AP and BQ to be two adjoining incident rays that follow PA' and QB', respectively, after reflecting on points P and Q on the Crystal surfaces. The highest value can be seen whenever the route variance between APA' and BQB' is an integral multiple of λ , indicating the presence of constructive interference.¹⁰

5.3. Advantages of X-ray diffraction crystallography

Strong and quick (less than 20 minutes) method for identifying an unidentified element.

In most cases, it provides an unambiguous mineral determination.

XRD units are widely available.

Data interpretation is relatively straightforward. Requires minimal sample preparation

Highly sensitive (the quality of being sensitive) Non corrosive, steady character/nature

Easy to prepare Sample/material Easy to handle or use

Cheap maintenance Work High/fast speed. Glancing angle.

Effective design.

Labour- saving system/ convenient. A system automatically operates.

Useful for identification of purity of the sample

5.4. Applications of X-ray crystallography

1. The primary application of X-ray powder diffraction is for the determination of unidentified mineral crystals (elements, inorganic chemicals, etc.).
2. Research findings in geology, environmental science, material science, engineering, and biology depend on determining the presence of unidentified substances.
3. Utilized to detect minerals with fine grains that are challenging to distinguish visually, such as clays and mixed layer clays.
4. To calculate the circumference of the unit cell.
5. For determining the crystal's fundamental composition.

6. It is employed to quantify textural features in a polycrystalline specimen, which includes grain orientation. By employing advanced methods, X-ray diffraction (XRD) may be utilized to quantify mineral quantities as well as identify the arrangement of crystals through Rietveld refinement.

5.5. To be particular about pharmaceutical field

Drugs come in a range of shapes and compositions, including tablets, pills, capsules, and aerosol sprays. An individual medicine can be manufactured in hundreds of approaches to improve its capacity to penetrate the circulatory system and deliver treatment to a particular area (skin's lotion, nasal spray, anti-itch, anticancer, etc.) and for it to function immediately or over time. A dosage form might increase the medication's shelf life, provide stomach buffering, enhance its flavour, or be a component of the plan for the drug's timed release into the bloodstream.

Pharmaceuticals' chemical makeup can be clearly identified using XRD. The crystal structures found in the medication under investigation directly influence the XRD pattern. As a result, it is easy to retrieve the parameters generally linked to crystal structure.

An X-ray powder diffraction pattern is required for examining the structure of the crystal, get a patent, and safeguard the investment made by the business. When it comes to multicomponent formulations, it is possible to accurately assess *in vivo* the proportions of the active ingredients in the final tablet forms as well as the proportion of the various amorphous packing materials used.¹²

XRD also possess various important application in different fields such as: -




Figure 2: Various application of XRD Instrumentation of XRD

5.6. Diffractometer

A diffractometer is a very accurate device with two independent rotational axes. With the aid of this apparatus, we are able to determine an X-ray beam's intensity data as a function of angle and to satisfy Bragg's law for X-rays with known wavelengths.¹³ The primary three elements of an X-

ray diffractometer are an X-ray tube, a sample holder, and an X-ray detector. By employing a filament that has been burned to create electrons, a voltage is applied to speed up the electrons towards a target, and the rapidly moving electrons attack the material of interest generating X-rays in a cathode ray tube. Once electrons possess enough force to remove the innermost electrons of the target material, characteristic X-ray spectra appear. Of the several components that comprise these spectrums, $K\alpha$ and $K\beta$ are particularly abundant.

Part of $K\alpha$ is composed of $K\alpha 1$ and $K\alpha 2$. Compared to $K\alpha 2$, $K\alpha 1$ has a slightly briefer spectrum and is twice as strong. The specific wavelengths (Cu, Fe, Mo, and Cr) correspond to the target material. Filtering with foils or crystal monochrometers needs to generate monochromatic X-rays for diffraction to take place. Since $K\alpha 1$ and $K\alpha 2$ have suitably near wavelengths, a weighted mean of them is used. Copper is a commonly frequently employed target material for single-crystal diffraction, with a Cu $K\alpha$ radiation of 1.5418

A. Focused X-rays will be applied to the specimen. When the specimen and sensor are spinning, the intensity of the reflecting X-rays is determined.

Once the object being photographed is impinged upon by the reflected X-rays' geometry, Bragg's law is achieved. This radiation from X-rays is detected and processed by a detector. After that, the data is transformed into a count rate and sent to a computer monitor or printer. The object being studied revolves at an angle μ in the course of the collimated X-ray beam in an X-ray diffractometer, while the X-ray detector turns at an angle of 2θ to collect the scattered X-rays. The tool employed to retain the position and spin the material is called a goniometer. In conventional particle patterns, statistics are collected at 2θ from 5 to 70°, angles which are set in the X-ray scanner.¹²

6. X-ray Diffraction Method

The x-ray diffraction behavior is helpful in determining the composition of solids. Additionally, for the X-ray spectrum investigation, Bragg's principle is often applied in such two scenarios for determining the crystal arrangement using Bragg's law¹⁰. Additional constant range of wavelengths, λ or θ , is given to enable to do this practically. In other words, given an amount of the inclination " θ ," the value of λ is randomly selected. In general, three approaches are used to examine the structure of molecules. These are the powder method, Laue method, and Rotating Crystal method.

6.1. Laue method

This is among the crucial techniques. Essentially employed to determine crystal symmetry, this tool is utilized for examining crystal structure.¹⁰




Figure 3: Laue method of XRD

In this procedure, a tiny crystal measuring 1 mm x 1 mm x 1 mm is put on a goniometer and exposed to a stream of polychromatic X-rays with frequencies that vary from 0.2 Å to 2 Å. The crystal's alignment with regard to the X-ray beam is capable of being altered by rotating the goniometer. Usually, the beam is let pass through parallel to the crystal's plane of analysis¹⁰.

When you traverse the crystal. The X-ray hits many Bragg's planes with d intervals between them and creating various angles " θ " with relation to the X-ray's incidence source. When λ and θ meet the Bragg's condition, then

$$2d \sin\theta = n\lambda$$

For a given value of d .

A diffraction pattern is produced by constructive interference and results in a rise in strength in particular planes. A

photographic plate placed on the contrary edge of the crystal will show this diffraction pattern. The pattern of diffraction found on the picture plate, comprises of a symmetrical spot layout based on the crystal lattice's symmetry characteristic. Typically, the degree of symmetry is found using the Laue technique.¹⁰

6.2. Rotating crystal method

Using this technique, a single, one-millimeter crystal is fixed to a revolving shaft, such that the spindle's rotational axis and one of the crystal's axis overlap. A monochromatic X-ray beam hits the substance parallel to the spindle's rotational axis.

A hollowed tubular container with an axis that is intersecting with the shaft's axis covers it, such a way that the crystal is positioned in the middle of the cylindrical holder, in order to acquire the pattern of diffraction. The surface of the cylindrical container has a photographic plate affixed to it. Note that the axis of motion is typically assumed to be the vertical direction.

Diffraction of x-rays will occur when the crystal rotates. Whenever the Crystalplane's position with relation to X rays, in a manner that satisfies Bragg's Condition. Every crystal

axis that is parallel to the spindle's rotational axis. The incident ray should be divided into a horizontal surface that is parallel to the shaft. On the other hand, the plane that includes the X-ray incident stream will not exhibit any scattering.¹⁰

6.3. Powder crystal method

One crystal has to be precisely mounted on one crystal axis in order to use Bragg's technique and the rotating Crystal technique. It is a laborious process to do. The powder crystal approach is utilized to get around this problem. Independent of one another, Deby, Scherer, and Hull devised this technique. The crystalline material is crushed into a powder using this process, which causes the crystallites to adopt a random orientation. A tiny capillary tube is filled with a sample of powdered powder, constructed of non-diffracting substance or, with the aid of non-diffracting binding substance, the sample is simply adhered to a hair. It is put in the line of a thin, monochromatic X-ray beam. Although there are additionally many crystallites with erratic orientations, the underlying premise of the crystal technique is this. For Small Angle X-ray Scattering (SAXS).

This approach explores the field of solution-based research, revealing the parameters and nature of macromolecules. SAXS was essential in exposing the constant modifications to the structure of proteins such as calmodulin in the analysis of protein conformational changes. Even though it might be difficult to precisely identify molecule shapes via scattered data, SAXS is nevertheless a useful technique for comprehending solution structures.¹⁴

7. Impactful Discoveries by Crystallography

The significance of X-ray diffraction and its uses in various research domains have resulted in the granting of 29 Nobel Prizes to date. The United Nations has recognized the effective introduction of crystallography into several scientific domains by designating 2014 as the International Year of Crystallography.¹⁵ Therefore, it is crucial to highlight a few groundbreaking findings that increased crystallography's significance.

A diamond's geometry made crystallography's role in understanding material characteristics clear after the first crystal structure of NaCl was identified by X-ray diffraction. A repeating quadrilateral arrangement of carbon atoms in the crystal lattice was found to be the cause of diamond's extreme hardness.¹⁶ Hexamethylenetetramine, quartz, and hex methylbenzene were among the next substances whose structures were determined. W.H. Barnes structural research on water were crucial in giving clear explanations of hydrogen bonding in water crystals in the late 1920s.¹⁷ Since that time, when it comes to examining the three-dimensional configuration of many other materials, such as elements,

Bragg's diffraction to occur, that is, for the diffraction to happen in line with the requirement.

$$2d \sin\theta = n\lambda,$$

Where λ is a constant in this instance, all potential diffraction planes must be accessible.

As a result, a collection of parallelograms that are angled at different angles towards the x-ray beam will produce the reflected image.¹⁰




Figure 4: Powder crystal method of

binary compounds, as well as proteins, X-ray diffraction has shown to be a trustworthy technique.

J. D. Bernal and colleagues gathered the first X-ray diffraction images of pepsin, a tiny hydrated protein, in 1934.¹⁸ About 25 years later, J. Kendrew et al. and M. Perutz et al. discovered the crystal structures of myoglobin and hemoglobin, respectively, and were honoured with the Nobel Prize in Chemistry in 1962. Many biological compounds, including insulin (1954), penicillin (1949), vitamin B12 (1954), and cholesterol (1937), were solved by D.C. Hodgkin (1969). In 1964, she received the Chemistry Nobel Prize in recognition of her ground-breaking works.

The finding of the blueprint for DNA by J. D. Watson and F. H. C. Crick, which was based on diffraction tests conducted by R. Franklin, was one of the most significant achievements of the twentieth century.¹⁹ Along with M. Wilkins, who had collaborated with Franklin, Watson, Crick, and the other recipient received the 1962 Nobel Prize in Physiology. The "double helix" discovery made macro molecules and protein crystallography possible, which are now vital resources in the biology and medical disciplines.²⁰

7.1. Need of X-ray diffraction and crystallography

A molecule with medicinal promise must pass several obstacles in the form of time, money, and human resources before it can be brought to market. This is a demanding procedure. It might take up to 20 years for a single chemical to be authorized as a medication, and research and development expenses can range from \$2 to \$3 billion.¹⁹ The scientific setting for drug development has changed between a focus on phenotypic identification or empirical assessment of

potential therapies to an emphasis on biological avenues in order to aid expedite the procedure and reduce the duration and effort needed for clinical testing and safety authorization.²¹

When combined alongside other identical techniques, X-ray structural analysis methods such as single crystal X-ray diffraction (SCXRD) and powder X-ray diffraction (PXRD) using the pair distribution function (PDF) have significantly aided in the conceptualization of biomimetic drugs alongside the continual assessment of drug stability. In the past, X-ray diffraction (XRD) has been the most popular method for identifying various structural characteristics in drug molecules. It has been helpful in the creation of specific biomolecule-ligand complexes, drug-receptor interactions, and structure-based drug design (SBDD).²²

The use of X-ray crystallography in drug design is crucial. The first step in developing a novel medicine to tackle a particular bacterium or virus is to identify a tiny chemical that can obstruct the enzymes and active proteins that are responsible for attacking human cells.

Scientists can create pharmacological molecules that can clamp onto the protein's "active" location and so inhibit its detrimental function by knowing the exact shape of the protein.⁸ With the advent of powerful X-ray sources (such as synchrotrons), novel detectors, innovative approaches to research, and state-of-the-art programmes, the number of accurately identified crystal structures of small and macromolecules has increased dramatically.

More than 750,000 small molecule crystal structures can currently be found in the Cambridge Structural Database (CSD)²³ and over 105,000 bio macromolecule structures can be found in the Protein Data Bank, approximately 90% of which were determined using X-ray diffraction.²⁴

Researchers could greatly improve X-ray crystallography by integrating AI approaches to more precisely and efficiently define macromolecular structures, find intriguing targets, and create novel medications. As a result, X-ray crystallography is well-positioned to continue playing a crucial part in discovering drugs as well as development into

the coming years.²

7.2. Pros of XRD when compared with other methods

Mass spectrometry, NMR spectroscopy, and X-ray crystallography have recently taken the lead in the structure investigation of natural compounds. From an instrumental standpoint, the MS field has made significant progress.²⁶ In the last ten years, new technologies including MSI, REIMS, IM-MS, and AS-MS have emerged. These resources, particularly when used in tandem, could provide natural product chemists new, standardized techniques.

Increased use of X-ray diffraction methods is anticipated. Conventional diffractometer's can determine the structure more easily from smaller particles or granulated specimens. In relation to speed, X-ray crystallography ought to potentially be better than NMR spectroscopy due to the growing availability of synchrotron sources worldwide and the falling costs of beamline time.

The full structure of a natural substance on the order of the size of taxol may now be revealed by renting 2 hours of synchrotron measurement time. These 120 minutes can cost between 2000 and 3000 EUR and cover data gathering, refinement, and evaluation. The same task might take one day on a typical laboratory diffractometer. On the other hand, the data collection and taxol interpretation for multidimensional NMR spectroscopy can require twice that time.²⁶

8. Protein structures may be atomically defined using X-ray diffraction (XRD), a key technique in structural biology. Since the discovery of the first protein structures, XRD has been crucial to understanding the molecular machinery of life. The accuracy and efficiency of XRD have significantly enhanced as science has made incremental progress. These advancements have been driven by the need to study tiny or more complex crystals, get higher-resolution frameworks, and conduct research more quickly. XRD is thus better than other techniques.²⁷

9. Conclusion

10. In pharmaceutical research, X-ray crystallography has shown to be a highly helpful tool. The advantages of using crystallography and x-ray diffraction are examined in this article, with an emphasis on how they are used in the pharmaceutical sector. Several x-ray diffraction techniques are theoretically examined, and the need of XRD is explained. A number of crystallographic findings are analyzed and evaluated.

References

Ummaleti R. XRD: Principle, Instrumentation, Types, and Uses. 2020; Nova Science Publishers.

2. Lee M. From the basics to the applications of X-ray diffraction in materials research. CRC Press, 1st ed., 2017, p. 302. 55 2025;12 (1):49–55 Parkhi et al./International Journal of Pharmaceutical Chemistry and Analysis

3. Tiwari S, Talreja M. S. Powder x-ray crystallography: an effective identification and analysis technique. *Mod Anal Int J Anal Experimen* 2020;12(12):109–112.

4. Martin J.H.J., Smyth M. S. X-ray crystallography. In 2000, *Mol Pathol.* 53(1): 8.

5. Applications of X-ray powder diffraction in drug screening and protein crystallography by Spiliopoulou M, Valmas A, Triandafyllidis D. P, Kosinas, C, Fitch, A, Karavassili F, and Margioliaki I. *Crystals.* 10(2):54, 2020.

6. Bragg WH. Bakerian Lecture: Crystal structure and X-rays. *Transactions of the Royal Society of London in Philosophy. Including Papers of a Physical or Mathematical Nature, Series A.*1915;215(523-37):253-274.

7. Suryawanshi R.P. and Kamat S. S. Using X-ray powder diffraction (XRPD) and inductively coupled plasma mass spectrometry (ICP-MS) in combination to create a herbal therapeutic product of BPC capsules. In 2017, *Eur J Biomed.* 4(02):360-6.

8. X-ray crystallography and its use in comprehending the physicochemical characteristics of medicinal cocrystals, Aidipapula S., & Vangala V. R. *Indian Journal of Inorganic Science,* 2017;97:227-43.

9. Das R, Ali E, and Hamid S. B. A Review of the Present Uses of X-Ray Powder Diffraction. 38:95-109 in *Rev Adv Mater Sci.* (2014).

10. Sharma, S. X-ray diffraction analysis: Fundamentals, tools, and applications. <https://physicswave.com/x-ray-diffraction-analysis-principle-instrument-and-applications/>, *Physics Wave,* 2020

11. Mogk, D., Goodge, J., Eby, N., and Henry, D. Bragg's Law with relation to X-ray reflection. *Geochemical Analysis and Instrumentation, 2017. Braggs Law.html* https://serc.carleton.edu/research_education/geochemsheets

12. X-ray diffraction: equipment and applications by Bunaciu A. A, Udrișteiu E.G, and Aboul-Enein H. Y. *Anal Chem. Crit Rev.* 2015; 45(4):289-99.

13. Waseda Y, Matsubara E, Shinoda K. *Introduction, Examples, and Solved Problems in X-ray Diffraction Crystallography.* 14. Asghar, Saira, Rehmat, Saira, and Iqtadar, Rabia. Springer Science & Business Media, 2011. An article reviewing the function of X-ray diffraction in the structural clarification of biological molecules. 2023; *Hamdard J Pharm.* 3:35-56.

In honor of the International Year of Crystallography, Desiraju G.R. *Design of Crystal Growth.* 2014;14(1):1-1.

Bragg WL, Bragg WH 16. The diamond's structure. *Proceedings of the London Royal Society. Papers of a Mathematical and Physical Character, Series A,* 1913;89(610):277-91.

17. Barnes, W. H. The crystal structure of ice between 0 C. and—183 C. *Proceedings of the London Royal Society. Series A, Containing Papers of a Mathematical and Physical Character.* 1929;125(799):670-93.

18. Bernal J.D, Crowfoot D., X-ray photographs of crystalline pepsin. *Nature*. 1934;133(3369):794-95.

19. Dabrowiak J. C. Metals in medicine. 2nd Edi. John Wiley & Sons. 2017.

20. Watson J.D, Crick F.H. Molecular structure of nucleic acids: a structure for deoxyribose nucleic acid. *Nature*. 1953;171(4356):737- 9.

21. Galkina Cleary E, Beierlein J.M, Khanuja N.S, McNamee L.M, Ledley F.D. Contribution of NIH funding to new drug approvals 2010–2016. *Proceed Nat Acad Sci*. 2018;115(10): 2329-34.

22. Rodríguez I, Gautam R, Tinoco A. D. Using X-ray diffraction techniques for biomimetic drug development, formulation, and polymorphic characterization. *Biomimetics*. 2020;6(1): 1.

23. Groom C.R, Bruno I.J, Lightfoot M.P, Ward S.C. The Cambridge structural database. *Structural Sci*. 2016;72(2): 171-9.

24. Berman H. M, Westbrook J, Feng Z, Gilliland G, Bhat T. N, Weissig H, & Bourne P. E. The protein data bank. *Nucl Acids Res*. 2000;28(1): 235-42.

25. Bijak V, Szczygiel M, Lenkiewicz J, et al. The current role and evolution of X-ray crystallography in drug discovery and development. *Expert Opin Drug Discov*. 2023;18(11):1221-30.

26. Přichystal J, Schug KA, Lemr K, Novák J, Havlíček V. Structural Analysis of Natural Products. *Anal Chem*. 2016;1:88(21):10338-46.

27. Smith, J., & Jones, A. Advances in X-ray Diffraction Techniques. *J Struc Biol*. 2023;214:12-24.